



Contents lists available at ScienceDirect

Saudi Journal of Biological Sciences

journal homepage: [www.sciencedirect.com](http://www.sciencedirect.com)

Original article

# Determination of experimental domain factors of polyphenols, phenolic acids and flavonoids of lemon (*Citrus limon*) peel using two-level factorial design

Zainol Haida<sup>a</sup>, Sharin Ab Ghani<sup>b</sup>, Jaafar Juhu Nakasha<sup>a</sup>, Mansor Hakiman<sup>a,\*</sup><sup>a</sup> Department of Crop Science, Faculty of Agriculture, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia<sup>b</sup> High Voltage Engineering Research Laboratory, Faculty of Electrical Engineering, Universiti Teknikal Malaysia, Melaka, Hang Tuah Jaya, 76100 Durian Tunggal, Melaka, Malaysia

## ARTICLE INFO

### Article history:

Received 22 November 2020

Revised 6 September 2021

Accepted 9 September 2021

Available online 16 September 2021

### Keyword:

*Citrus limon*

Polyphenols

Phenolic acids

Flavonoids

Two-level factorial design

## ABSTRACT

This study aimed to evaluate the significant extraction factors in achieving higher recovery yield of total polyphenols, phenolic acids and flavonoids content from *Citrus limon* peel using two-level factorial design. The effect of five independent factors including drying temperature (40–60 °C), methanol concentration (20–60%), extraction temperature (28–60 °C), extraction time (30–60 min) and storage duration (0–14 days) were evaluated. Among all the examined factors, results showed that drying temperature, storage duration and extraction temperature were the most significant and contributing factors affecting the total polyphenols, phenolic acids and flavonoids content of lemon peel at  $P < 0.05$ . On the contrary, methanol concentration and extraction time exhibited the least significant and contribution at  $P$  greater than 0.05. In conclusion, the experimental domain factors were successfully obtained from this experiment. Therefore, further study on optimization of the obtained factors will be conducted in the future study using response surface methodology.

© 2021 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

*Citrus limon* or commercially known as lemon belongs to the family Rutaceae. Lemon is one of the most popular *Citrus* species in the world after orange and mandarin (Miran et al., 2016). In other countries, lemon is known as zitrone in Germany, le citron in French, limon in Spanish and ningmeng in Chinese (Klimek-Szczykutowicz et al., 2020). Lemon is popular due to its various types of phytochemical and ample supply of vitamin C, folic acid, potassium and pectin which act as a supplement to improve human health and prevent diseases (Rafiq et al., 2018; Sagar et al., 2018). Lemon is widely consumed worldwide as a fresh fruit and juice but the lemon peel is mostly discarded as a waste (Sagar et al., 2018). However, consumers are not aware that the lemon peel contains a

wide range of bioactive compounds with substantial antioxidant activity which might contain useful bioactive compounds comparable to that of lemon pulp (Mahato et al., 2019). The lemon peel can be divided into epicarp or flavedo (colored peripheral surface) and mesocarp or albedo (white soft middle layer) (Rafiq et al., 2018). Since the lemon peel is considered as a waste, only several studies have been conducted on phenolics content and antioxidant activities of lemon peel. In order to maximize the phenolics content and antioxidant activities of lemon peel, optimization of the extraction process is an important step that needs to be carried out properly.

The optimization of the extraction process is a crucial step as different species or plant parts require distinct extraction procedures and conditions to yield maximum phenolic compounds and antioxidant activities due to their different characteristics and phytochemical constituents (Azwanida et al., 2015). In the optimization of phenolics content extraction from *Citrus* species, the frequently used extraction variables are solvent concentration, liquid–solid ratio, extraction temperature and extraction time and the ranges of each variable used are varied depending on several factors including plant parts, maturity of fruits and targeted secondary metabolites (Assefa et al., 2017; Nipornram et al., 2018; Fakayode and Abobi, 2018; Colodel et al., 2018; Nishad et al., 2019; Kim and Lim, 2020).

\* Corresponding author at: Department of Crop Science, Faculty of Agriculture, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia.

E-mail address: [mhakiman@upm.edu.my](mailto:mhakiman@upm.edu.my) (M. Hakiman).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

In this current study, two-level factorial design was adopted as a statistical screening process. The two-level factorial design is a useful method for the extraction process to determine the main factor as well as interaction between the factors with minimal experimental runs. Hence, this study was conducted to screen the independent factors including drying temperature, methanol concentration, extraction temperature, extraction time and storage duration for the recovery of total polyphenols, phenolic acids and flavonoids content from lemon peel using two-level factorial design.

## 2. Materials and methods

### 2.1. Plant material

Fresh fruits of *Citrus limon* (lemon) at their mature stage were purchased from local grocery store in Selangor, Malaysia. The fruits were imported from South Africa. The fruits were cleaned thoroughly under running tap water. The fruit peel was manually separated from the flesh and wrapped with paper for drying procedure.

### 2.2. Chemicals and reagents

Gallic acid, rutin, Folin-Ciocalteu reagent, aluminium chloride, sodium carbonate and methanol were purchased from R&M chemical company and Sigma-Aldrich. All chemicals and reagents used in this study were of analytical grade.

### 2.3. Extraction process

The extraction procedure of lemon peel was carried out using maceration method. First, the dried samples which were dried at different oven temperatures (40 – 60 °C) were finely ground using a commercial blender. A total of 0.5 g of powdered sample was mixed with 25 mL of methanol at different concentrations (20 – 60%) as extraction solvent. Then, the mixtures were placed in a water bath with a temperature ranged from 28 to 60 °C and extraction time was set between 30 and 60 min. The mixtures were filtered using filter paper No. 1 and stored in the refrigerator (5 °C) for 0 to 14 days. The variables were set based on the design matrix generated by Design Expert software version 11.0 (Table 1).

### 2.4. Total polyphenols content

The method described by Marinova et al. (2005) was used for total polyphenols content analysis. Briefly, the extract (50 µL) was mixed with ten-fold diluted Folin-Ciocalteu reagent (1.25 mL) and the mixture was incubated for five minutes. After the incubation, 7% sodium carbonate (1.25 mL) added and the mixture was incubated at room temperature for an hour. The change in absorbance was measured using UV-Vis spectrophotometer at 725 nm. The gallic acid was used as standard and the total

polyphenols content of lemon peel was expressed as mg gallic acid equivalent per gram dry weight of sample (mg GAE/g DW).

### 2.5. Total phenolic acids content

The quantification of total phenolic acids content was conducted using method as explained by Singleton and Rossi (1965) and Haida et al. (2019). The mixture containing extract (0.5 mL), distilled water (4.5 mL) and Folin-Ciocalteu reagent (0.5 mL) were mixed and incubated for five minutes. After the incubation period, 7% sodium carbonate (5 mL) and distilled water (2 mL) were added. The reaction mixture was incubated at room temperature for 90 min and absorbance was measured at 750 nm using UV-Vis spectrophotometer. The gallic acid was used as standard and the total phenolic acids content of lemon peel was expressed as mg gallic acid equivalent per gram dry weight of sample (mg GAE/g DW).

### 2.6. Total flavonoids content

The determination of total flavonoids content from lemon peel was quantified using aluminium chloride colorimetric method as described by Marinova et al. (2005). The mixture of extract (0.5 mL), distilled water (2 mL) and 5% sodium nitrite (150 µL) were mixed thoroughly and incubated for five minutes. Then, 10% aluminium chloride (150 µL) was added to the mixture followed by 1 M sodium hydroxide (1 mL) and distilled water (1.2 mL). The reaction mixture was mixed and absorbance was measured at 510 nm using UV-Vis spectrophotometer. The rutin was used as standard and the total flavonoids content was expressed as mg rutin equivalent per gram dry weight of sample (mg RE/g DW).

### 2.7. Experimental design and statistical analysis

In this study, five extraction factorials namely drying temperature, methanol concentration, extraction temperature, extraction time and storage duration were analyzed using two-level factorial design. Each factorial including drying temperature (A, °C), methanol concentration (B, %), extraction temperature (C, °C), extraction time (D, minutes) and storage duration (days) were evaluated using two coded levels in a randomized trend with a total of 32 experimental trials ( $2^5 = 32$ ) and three replications of the centre points (Table 2). Three responses recorded were total polyphenol content (mg GAE/g DW), total phenolic acids content (mg GAE/g DW) and total flavonoids content (mg RE/g DW). The experimental runs and analysis were conducted in triplicate and the value was expressed as mean. The analysis of variance (ANOVA) was used to analyzed the p-value and significance of the model. The Design Expert 11.0 software (Stat-Ease Inc., Minneapolis, USA) was used for designing the experiment and data analysis.

**Table 1**  
Parameters for the two-level factorial design.

Factor	Notation	Factor levels		
		Low (-1)	Centre point (0)	High (+1)
Drying temperature (°C)	A	40	50	60
Methanol concentration (%)	B	20	40	60
Extraction temperature (°C)	C	28	44	60
Extraction time (min)	D	30	45	60
Storage duration (day)	E	0	7	14

**Table 2**  
Design matrix and the responses of two-level factorial design.

Run	Factors					Responses		
	A	B	C	D	E	Y <sub>1</sub>	Y <sub>2</sub>	Y <sub>3</sub>
1	60	20	60	60	14	2.8	4.74	47.26
2	60	60	60	60	14	3.08	5.05	55.7
3	60	20	28	30	0	2.5	7.9	37.4
4	60	60	28	30	0	2.4	8.16	37.12
5	60	20	28	30	14	2.78	6.05	42.37
6	60	60	60	30	0	2.77	8.04	47.12
7	40	20	28	60	0	1.86	3.89	21.51
8	40	20	60	60	0	1.94	3.48	28.45
9	60	60	28	30	14	2.9	4.74	31.87
10	60	60	28	60	0	2.58	7.22	35.9
11	40	60	60	30	0	2.71	5.12	37.4
12	40	20	60	30	0	2.07	4.86	28.54
13	40	20	28	60	14	1.82	3.55	22.79
14	60	20	60	60	0	2.52	8.11	45.26
15	60	60	60	30	14	3.24	5.62	54.29
16	40	60	60	60	14	3.55	6.97	54.31
17	40	60	28	30	0	1.72	3.77	22.4
18	60	60	28	60	14	2.92	4.54	38.76
19	40	60	28	60	0	1.4	3.71	21.31
20	60	20	60	30	14	2.92	6.06	49.45
21	60	60	60	60	0	2.78	6.78	47.01
22	60	20	28	60	14	2.71	4.94	44.76
23 <sup>a</sup>	50	40	44	45	7	2.58	4.52	34.04
24	60	20	28	60	0	2.45	7.42	32.34
25	40	60	60	60	0	2.62	5.24	33.62
26 <sup>a</sup>	50	40	44	45	7	2.69	4.33	34.7
27	40	20	28	30	0	1.96	5.55	22.26
28	40	60	28	60	14	1.99	4.11	23.93
29	40	20	60	30	14	2.24	4.31	30.93
30	40	20	28	30	14	1.79	3.68	24.01
31	60	20	60	30	0	2.63	8.87	41.43
32	40	60	60	30	14	2.74	5.17	38.84
33 <sup>a</sup>	50	40	44	45	7	2.65	4.2	34.45
34	40	20	60	60	14	2.29	5.12	34.54
35	40	60	28	30	14	2.06	4.18	22.7

A – Drying temperature (°C); B – Methanol concentration (%); C – Extraction temperature (°C); D – Extraction time (min); E – Storage duration (day); Y<sub>1</sub> – Total polyphenols content (mg GAE/g DW); Y<sub>2</sub> – Total phenolic acids content (mg GAE/g DW); Y<sub>3</sub> – Total flavonoids content (mg RE/g DW). <sup>a</sup>Centre point.

### 3. Results

#### 3.1. Influence of extraction factors on the total polyphenols content

The response model was highly significant with P < 0.0001 (Table 3). Based on Fig. 1 and Table 3, it can be observed that an increase in the drying temperature, methanol concentration, extraction temperature and storage duration resulted in higher recoveries of total polyphenols content of lemon peel. In contrary, extraction time showed the least effect on total polyphenols con-

**Table 3**  
P-value of factors on responses.


Factors	p-values for the responses		
	Y <sub>1</sub>	Y <sub>2</sub>	Y <sub>3</sub>
A	<0.0001	<0.0001	<0.0001
B	0.0007	0.9758	0.0227
C	<0.0001	0.0097	<0.0001
D	0.9051	0.0567	0.3489
E	0.0001	<0.0001	0.0007
Model	<0.0001	<0.0001	<0.0001
Adjusted R <sup>2</sup>	0.8396	0.8273	0.8988
C.V. (%)	7.72	11.70	9.09

A – Drying temperature (°C); B – Methanol concentration (%); C – Extraction temperature (°C); D – Extraction time (min); E – Storage duration (day); Y<sub>1</sub> – Total polyphenols content (mg GAE/g DW); Y<sub>2</sub> – Total phenolic acids content (mg GAE/g DW); Y<sub>3</sub> – Total flavonoids content (mg RE/g DW).

tent of lemon peel. The recovery of total polyphenols was greatly affected by drying temperature, methanol concentration, extraction temperature and storage duration with P < 0.05. Meanwhile, the extraction time was non-significant on total phenolic acids content. The effect of drying temperature was the major contributing factor with 34.91%, followed by extraction temperature (20.47%), storage duration (9.94%) and extraction temperature (7.17%), respectively (Table 4). The Pareto Chart is used to measure the sampling error of individual variable and their interaction through the standard deviation. Based on Pareto chart of total polyphenols content (Fig. 1a), the t-value limit is 2.08596. Any factor or interaction that falls below t-value will be non-significant. In Fig. 1a, the Pareto chart showed that drying temperature was the most contributing factor in producing the highest total polyphenols content from lemon peel. In contrast, extraction time was not significant as it fell below t-value limit.

#### 3.2. Influence of extraction factors on the total phenolic acids content

For the total phenolics content, the response model was highly significant with P < 0.0001 (Table 3). Based on the Fig. 2, as the drying temperature and extraction temperature increased, the recoveries of total phenolic acids content were increased. For the storage duration, increased in incubation time resulted in lower recovery while methanol concentration and extraction time showed the least effect on total phenolic acids content of lemon peel (Table 3, Fig. 2). The recovery of total phenolic acids content



**Fig. 1.** (a) The Pareto Chart of all factors and interactions on total polyphenols content from lemon peel; Influence of (b) drying temperature; (c) methanol concentration; (d) extraction temperature; (e) extraction time; (f) storage duration on the total polyphenols content recovery from lemon peel. A greater slope shows a greater influence on the recovery.

was greatly affected by drying temperature, extraction temperature and storage duration with  $P < 0.05$ . Meanwhile, the factors of methanol concentration and extraction time were showed non-significant effect on total phenolic acids content. Based on Table 4, the major contributing factor of total phenolic acids content was drying temperature with 38.39%, followed by storage duration (14.37%) and extraction temperature (3.96%), respectively. Based on Pareto chart (Fig. 2a), drying temperature was the most contributing factor in extracting higher phenolic acids from lemon peel. Meanwhile, methanol concentration was non-significant as it fell below t-value limit (2.0639).

### 3.3. Influence of extraction factors on the total flavonoids content

In the analysis of total flavonoids content of lemon peel, the response model was highly significant with  $P < 0.0001$  (Table 3). Based on the Table 3 and Fig. 3, the recovery of total flavonoids content was increased as the drying temperature, methanol concentration, extraction temperature and storage duration were increased. Meanwhile, extraction time exhibited the lowest effect on total flavonoids content. The recovery for total flavonoids content of lemon peel was highly affected by drying temperature, methanol concentration, extraction temperature and storage dura-

**Table 4**  
Percentage contribution of factors on responses.

Factors	Percentage contribution (%)		
	Y <sub>1</sub>	Y <sub>2</sub>	Y <sub>3</sub>
A	34.91	38.39	43.55
B	7.17	4.67 × 10 <sup>-4</sup>	2.15
C	20.47	3.96	33.27
D	5.91 × 10 <sup>-3</sup>	2.01	0.33
E	9.94	14.37	5.37
AB	0.88	2.33	1.03
AC	6.77	1.18	0.31
AD	0.09	1.42	0.05
AE	0.11	19.08	0.02
BC	6.50	0.96	5.16
BD	0.32	0.91	0.30
BE	1.78	0.61	1.4 × 10 <sup>-4</sup>
CD	0.17	0.17	0.26
CE	0.21	0.74	1.13
DE	0.50	1.22	1.15

A – Drying temperature (°C); B – Methanol concentration (%); C – Extraction temperature (°C); D – Extraction time (min); E – Storage duration (day); Y<sub>1</sub> – Total polyphenols content (mg GAE/g DW); Y<sub>2</sub> – Total phenolic acids content (mg GAE/g DW); Y<sub>3</sub> – Total flavonoids content (mg RE/g DW).

tion with  $P < 0.05$ . Besides that, non-significant effect was shown by extraction time on total flavonoids content. Based on Table 4, drying temperature was the major contributing factor with 43.55%, followed by extraction temperature (33.27%), storage duration (5.37%) and methanol concentration (2.14%), respectively. Based on Pareto chart in Fig. 3a, drying temperature was the highest contributing factor in attaining highest total flavonoids content from lemon peel. Besides that, extraction time was not significantly affected the total flavonoids content as it fell below t-value limit (2.05553).

#### 4. Discussion


Among all the factors, drying temperature has the greatest influence on total polyphenols, phenolic acids and flavonoids content of lemon peel. From Table 3 and 4, this factors showed a highly significant value with  $P < 0.0001$  and it showed the highest percentage contribution for all the responses. Similarly, by comparing the line plots shown in Fig. 1b, 2b and 3b, the effect of drying temperature was clearly shown the greatest gradient of the slope. The total polyphenols, phenolic acids and flavonoids content extracted from the lemon peel was significantly increased as the drying temperature increased from 40 to 60 °C. The trend in our study is similar to previously published results *Solanum lycopersicum* (Gumusay et al., 2015; Azeez et al., 2019), leafy vegetables (Nwozo et al., 2015) and red rice (Hu et al., 2017) which observed the phenolics content was increased as the drying temperature increased. The accumulation of high phenolics during high drying temperature is related to the increment of the release phenolics bound from the cell wall as a result of ester breaks down between phenolic and cell wall due to the heat treatment (Azeez et al., 2019). As a result, more phenolics would be extracted from the sample.

Extraction temperature also plays an important role in extracting high total polyphenols, phenolic acids and flavonoids content. This factor exhibited highly significant effect on total polyphenols and flavonoids content with  $P < 0.0001$  and significant effect on total phenolic acids content with  $P < 0.05$ , respectively (Table 3). Based on Table 4, this factor also showed high percentage of contribution on total polyphenols, phenolic acids and flavonoids content and Fig. 1d, 2d and 3d showed large gradient of the slope. In this study, increment of extraction temperature was significantly increased the total polyphenols, phenolic acids and flavonoids

content of lemon peel. The possibilities suggested for the increment of phenolics content might be due to release of high amounts of phenolics due to the thermal destruction of cell walls and sub-cellular compartments. The thermal treatments during extraction process will soften the plant biomass which resulted in higher bioavailability of polyphenols content inside the cell wall. Besides that, heat can break the supramolecular structure of the plant biomass which release more phenolics. As a result, phenolic solubility, diffusion rate and mass transfer are increased while solvent viscosity and surface tension are decreased (Bunea et al., 2008; Nwozo et al., 2015; Mokrani and Madani, 2016). The effect of extraction temperature on phenolic content also have been reported by several previous studies on different types of plant sample (Thoo et al., 2010; Deng et al., 2017; Zhou et al., 2018).

It was also noticeable that storage duration was important in attaining high total polyphenols, phenolic acids and flavonoids content from lemon peel. From Table 3, storage duration was highly significant on total phenolic acids content with  $P < 0.0001$  and significant P-value was obtained on total polyphenols and flavonoids content with  $P < 0.05$ . From the line plot in Fig. 2f, negative slope gradient was obtained on total phenolic acids content, contrary with total polyphenols and flavonoids content which were produced the positive slope gradient (Fig. 1f and 3f). This showed that the phenolic acids content was decreased as the storage duration increased. However, the total polyphenols and flavonoids were increased as the storage duration was increased in this study. Polyphenols is one of the secondary metabolites that is known to be unstable during storage especially at ambient temperatures (Howard et al., 2012). However, at low temperature ( $<10$  °C) storage condition, the phenolics and flavonoids in fruits can be retained for several days. The study conducted by Fawole and Opara (2013) on pomegranate found that the total phenolics content was relatively stable for four-week storage duration at a temperature of 5, 7 and 10 °C. The deterioration of total phenolics content was observed at all temperatures after eight-week of storage duration (Fawole and Opara, 2013). The accumulation and deterioration of phenolics during storage as response to storage temperature and duration are mainly affected the genes and enzymes synthesis which is different in each type of food (Galani et al., 2017).

The extraction solvent is one of the factors that highly influenced the extraction of phenolics from plant sample. High polarity of solvent such as aqueous, methanol and ethanol are frequently used as extraction solvent (Dai and Mumper, 2010). In this current study, methanol was chosen to extract the phenolics content from the lemon peel. The uses of methanol as extraction solvent are highly preferred and previously reported this solvent is effective for phenolics extraction (Ye et al., 2015; Vajić et al., 2015). The results in this study found that methanol concentration was significantly affected the total polyphenols and flavonoids content extracted from lemon peel with  $P < 0.05$  (Table 3). However, methanol concentration was not significantly affected the total phenolic acids content with the P-value recorded was more than 0.05. Based on Fig. 1c and 3c, only small gradient slope was obtained on total polyphenols and flavonoids content. In term of percentage of contribution, total phenolic acids content recorded the lowest percentage of contribution than total polyphenols and flavonoids content (Table 4). The total polyphenols and flavonoids content were significantly increased as the methanol concentration increased from 20 to 60%. In contrast with the total polyphenols and flavonoids content, the increment of methanol concentration was not significantly increased the phenolic acids content. In plant, the polyphenols are present at different affinities indicating that different polarities of solvent might directly affecting the phenolics content extraction (Viera et al., 2017). Thus, this study showed that high concentration of solvent produced higher




**Fig. 2.** (a) The Pareto Chart of all factors and interactions on total phenolic acids content from lemon peel; Influence of (b) drying temperature; (c) methanol concentration; (d) extraction temperature; (e) extraction time; (f) storage duration on the total phenolic acids content recovery from lemon peel. A greater slope shows a greater influence on the recovery.

polyphenols and flavonoids content. But, a higher concentration of solvent approximately 80% of solvent is needed to extract higher amount of phenolic acids. According to Chirinos et al. (2007), low phenolics concentration at low concentration of extraction solvent may be due to high amount of water in solvent which is responsible to extract impurities (polysaccharides, organic acids and carbohydrates) may interfere the phenolics extraction.

Extraction time did not show any significant effect on total polyphenols, phenolic acids and flavonoids content in comparison

with other factors. The P-value recorded on all responses was more than 0.05 (Table 3). In addition, the percentage contribution recorded also was low (Table 4). In this study, the extraction time was conducted in 30 to 60 min. The non-significant results obtained in all responses might be due to the extraction time was too short. This phenomenon is explained by Fick's second law of diffusion, predicting that a final equilibrium between the solute concentrations in the solid matrix (plant matrix) and in the bulk solution (solvent) might be reached after a certain time



**Fig. 3.** (a) The Pareto Chart of all factors and interactions on total flavonoids content from lemon peel; Influence of (b) drying temperature; (c) methanol concentration; (d) extraction temperature; (e) extraction time; (f) storage duration on the total flavonoids content recovery from lemon peel. A greater slope shows a greater influence on the recovery.

(Silva et al., 2007). Moreover, prolongation of extraction time until reach the maximum limit will increase the chance of sample to denature. Thus, higher phenolics content can be extracted from the sample. In a previous study on *Citrus aurantifolia*, *Citrus nobilis*, *Citrus grandis* and *Citrus reticulata*, the optimum extraction time recorded was between 80 and 120 min (Tran et al., 2019; Hien et al., 2018; Colodel et al., 2018; Dao et al., 2020). Hence, it is suggested that the extraction time needed for maximum phenolic extraction from *Citrus limon* is more than 60 min.

**5. Conclusions**

The two-level factorial design was adopted in screening of five independent factors including drying temperature, methanol concentration, extraction temperature, extraction time and storage duration of lemon peel extraction. In attaining high total polyphenols, phenolic acids and flavonoids content from lemon peel extract, each factor was evaluated to find the most significant with high contribution factor. Results indicated that drying tempera-

ture, storage duration and extraction temperature were the most significant and contributing factor affecting the total polyphenols, phenolic acids and flavonoids content of lemon peel with  $P$ -value $<0.05$ . Meanwhile, extraction time exhibited the least significance and contribution with  $P>0.05$ . Therefore, optimization using the obtained significant factors will be conducted using response surface methodology in the future.

### CRedit authorship contribution statement

**Zainol Haida:** Conceptualization, Methodology, Investigation, Data curation, Software, Formal analysis, Writing – original draft and editing the manuscript. **Sharin Ab Ghani:** Conceptualization, Methodology, Software, Formal analysis, Writing – review and editing. **Jaafar Juju Nakasha:** Project administration, Supervision, Funding acquisition, Resources, Writing – review and editing. **Man-sor Hakiman:** Project administration, Supervision, Funding acquisition, Resources, Writing – review and editing. All authors have made direct contribution and this manuscript was approved by all authors for publication.

### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Acknowledgements

Authors would like to thank the Ministry of Education Malaysia and Universiti Putra Malaysia for financing the present study through Translational Research Program, 5526700, Transdisciplinary Research Grant Scheme, 5535802 and Putra Grant – Putra Young Initiative, 9629000.

### Author contributions

Z.H. was contributed in conceptualization, methodology, investigation, data curation, software, formal analysis, writing – original draft and editing the manuscript. S. A. G. was conceptualization, methodology, software, formal analysis and writing – review and editing. J. J. N. and M. H. were contributed in project administration, supervision, funding acquisition, resources and writing – review and editing. All authors have made direct contribution and this manuscript was approved by all authors for publication.

### References

Assefa, A.D., Saini, R.K., Keum, Y.S., 2017. Extraction of antioxidants and flavonoids from yuzu (*Citrus junos* Sieb ex Tanaka) peels: A response surface methodology study. *J. Food Meas. Charact.* 11 (2), 364–379.

Azwanida, N.N., 2015. A review on the extraction methods use in medicinal plants, principle, strength and limitation. *Med. Aromat. Plants*, 4, 2167–210412.

Azeez, L., Adebisi, S.A., Oyediji, A.O., Adetoro, R.O., Tijani, K.O., 2019. Bioactive compounds' contents, drying kinetics and mathematical modelling of tomato slices influenced by drying temperatures and time. *J. Saudi Soc. Agric. Sci.* 18 (2), 120–126.

Bunea, A., Andjelkovic, M., Socaciu, C., Bobis, O., Neacsu, M., Verhé, R., Camp, J.V., 2008. Total and individual carotenoids and phenolic acids content in fresh, refrigerated and processed spinach (*Spinacia oleracea* L.). *Food Chem.* 108 (2), 649–656.

Chirinos, R., Rogez, H., Campos, D., Pedreschi, R., Larondelle, Y., 2007. Optimization of extraction conditions of antioxidant phenolic compounds from mashua (*Tropaeolum tuberosum* Ruiz & Pavón) tubers. *Sep. Purif. Technol.* 55 (2), 217–225.

Colodel, C., Vriesmann, L.C., Teófilo, R.F., de Oliveira Petkowicz, C.L., 2018. Extraction of pectin from ponkan (*Citrus reticulata* Blanco cv. Ponkan) peel: Optimization and structural characterization. *Int. J. Biol. Macromol.* 117, 385–391.

Dai, J., Mumper, R.J., 2010. Plant phenolics: Extraction, analysis and their antioxidant and anticancer properties. *Molecules* 15 (10), 7313–7352.

Dao, T.P., Tran, T.H., Nguyen, P.T.N., Tran, T.K.N., Ngo, T.C.Q., Nhan, L.T.H., Anh, T.T., Toan, T.Q., Quan, P.M., Linh, H.T.K., 2020. Optimization of microwave assisted hydrodistillation of essential oil from lemon (*Citrus aurantifolia*) leaves: Response surface methodology studies. In *IOP Conference Series: Materials Science and Engineering* (Vol. 736, No. 2, p. 022038). IOP Publishing.

Deng, J., Xu, Z., Xiang, C., Liu, J., Zhou, L., Li, T., Yang, Z., Ding, C., 2017. Comparative evaluation of maceration and ultrasonic-assisted extraction of phenolic compounds from fresh olives. *Ultrason. Sonochem.* 37, 328–334.

Fakayode, O.A., Abobi, K.E., 2018. Optimization of oil and pectin extraction from orange (*Citrus sinensis*) peels: A response surface approach. *J. Anal. Sci. Tech.* 9 (1), 1–16.

Fawole, O.A., Opara, U.L., 2013. Effects of storage temperature and duration on physiological responses of pomegranate fruit. *Ind. Crops Prod.* 47, 300–309.

Galani, J.H., Patel, J.S., Patel, N.J., Talati, J.G., 2017. Storage of fruits and vegetables in refrigerator increases their phenolic acids but decreases the total phenolics, anthocyanins and vitamin C with subsequent loss of their antioxidant capacity. *Antioxidants*. 6 (3), 59.

Gümüsay, Ö.A., Borazan, A.A., Ercal, N., Demirkol, O., 2015. Drying effects on the antioxidant properties of tomatoes and ginger. *Food Chem.* 173, 156–162.

Haida, Z., Syahida, A., Ariff, S.M., Maziah, M., Hakiman, M., 2019. Factors affecting cell biomass and flavonoid production of *Ficus deltoidea* var. kunstleri in cell suspension culture system. *Sci. Rep.* 9 (1), 1–8.

Hien, T.T., Nhan, N.P.T., Trinh, N.D., Ho, V.T.T., Bach, L.G., 2018. Optimizing the pomelo oils extraction process by microwave-assisted hydro-distillation using soft computing approaches. *Solid State Phenom.* 279, 217–221.

Howard, L.R., Prior, R.L., Liyanage, R., Lay, J.O., 2012. Processing and storage effect on berry polyphenols: Challenges and implications for bioactive properties. *J. Agric. Food Chem.* 60 (27), 6678–6693.

Hu, Z., Tang, X., Liu, J., Zhu, Z., Shao, Y., 2017. Effect of parboiling on phytochemical content, antioxidant activity and physicochemical properties of germinated red rice. *Food Chem.* 214, 285–292.

Kim, D.S., Lim, S.B., 2020. Extraction of flavanones from immature *Citrus unshiu* pomace: Process optimization and antioxidant evaluation. *Sci. Rep.* 10 (1), 1–13.

Klimek-Szczykutowicz, M., Szopa, A., Ekiert, H., 2020. *Citrus limon* (Lemon) phenomenon—A review of the chemistry, pharmacological properties, applications in the modern pharmaceutical, food, and cosmetics industries, and biotechnological studies. *Plants*. 9, 119.

Mahato, N., Sinha, M., Sharma, K., Koteswararao, R., Cho, M.H., 2019. Modern extraction and purification techniques for obtaining high purity food-grade bioactive compounds and value-added co-products from citrus wastes.  *Foods*. 8, 523.

Marinova, D., Ribarova, F., Atanassova, M., 2005. Total phenolics and total flavonoids in Bulgarian fruits and vegetables. *J. Uni. Chem. Tech. Metal.* 40, 255–260.

Miran, W., Nawaz, M., Jang, J., Lee, D.S., 2016. Sustainable electricity generation by biodegradation of low-cost lemon peel biomass in a dual chamber microbial fuel cell. *Int. Biodeterior. Biodegradation.* 106, 75–79.

Mokrani, A., Madani, K., 2016. Effect of solvent, time and temperature on the extraction of phenolic compounds and antioxidant capacity of peach (*Prunus persica* L.) fruit. *Sep. Pur. Tech.* 162, 68–76.

Nipornram, S., Tochampa, W., Rattanatraiwong, P., Singanusong, R., 2018. Optimization of low power ultrasound-assisted extraction of phenolic compounds from mandarin (*Citrus reticulata* Blanco cv. Sainampung) peel. *Food Chem.* 241, 338–345.

Nishad, J., Saha, S., Dubey, A.K., Varghese, E., Kaur, C., 2019. Optimization and comparison of non-conventional extraction technologies for *Citrus paradisi* L. peels: A valorization approach. *J. Food Sci. Tech.* 56 (3), 1221–1233.

Nwozo, S.O., Oso, B.J., Oyinloye, B.E., 2015. Effect of heat on antioxidant activity of some tropical leafy vegetables. *Nig. J. Basic Appl. Sci.* 23 (2), 93–101.

Rafiq, S., Kaul, R., Sofi, S.A., Bashir, N., Nazir, F., Nayik, G.A., 2018. Citrus peel as a source of functional ingredient: A review. *J. Saudi Soc. Agric. Sci.* 17, 351–358.

Sagar, N.A., Pareek, S., Sharma, S., Yahia, E.M., Lobo, M.G., 2018. Fruit and vegetable waste: Bioactive compounds, their extraction, and possible utilization. *Compr. Rev. Food Sci. Food Saf.* 17 (3), 512–531.

Silva, Souza, Rogez, Rees, Larondelle, 2007. Antioxidant activities and polyphenolic contents of fifteen selected plant species from the Amazonian region. *Food Chemistry* 101 (3), 1012–1018. <https://doi.org/10.1016/j.foodchem.2006.02.055>. In this issue.

Singleton, V.L., Rossi, J.A., 1965. Colorimetry of total phenolics with phosphomolybdc-phosphotungstic acid reagent. *Am. J. Enolo. Viticult.* 16, 144–158.

Thoo, Y.Y., Ho, S.K., Liang, J.Y., Ho, C.W., Tan, C.P., 2010. Effects of binary solvent extraction system, extraction time and extraction temperature on phenolic antioxidants and antioxidant capacity from mengkudu (*Morinda citrifolia*). *Food Chem.* 120 (1), 290–295.

Tran, T.H., Nguyen, P.T.N., Ho, V.T.T., Le, T.H.N., Bach, L.G., Nguyen, T.D., 2019. Using soft computing approaches for orange (*Citrus nobilis* Lour. var. *nobilis*) oils extraction process. *Mater. Sci. Eng.* 479, 012015. <https://doi.org/10.1088/1757-899X/479/1/012015>.

Vajić, U.-J., Grujić-Milanović, J., Živković, J., Šavikin, K., Godevac, D., Miloradović, Z., Bugarski, B., Mihailović-Stanojević, N., 2015. Optimization of extraction of stinging nettle leaf phenolic compounds using response surface methodology. *Ind. Crops Prod.* 74, 912–917.

Viera, V.B., Piovesan, N., Rodrigues, J.B., de O Mello, R., Prestes, R.C., dos Santos, R.C. V., de A Vaucher, R., Hautrive, T.P. and Kubota, E.H., 2017. Extraction of phenolic



- compounds and evaluation of the antioxidant and antimicrobial capacity of red onion skin (*Allium cepa* L.). *Int. Food Res. J.* 24(3), 990–999.
- Ye, F., Liang, Q., Li, H., Zhao, G., 2015. Solvent effects on phenolic content, composition, and antioxidant activity of extracts from florets of sunflower (*Helianthus annuus* L.). *Ind. Crops Prod.* 76, 574–581.
- Zhou, P., Wang, X., Liu, P., Huang, J., Wang, C., Pan, M., Kuang, Z., 2018. Enhanced phenolic compounds extraction from *Morus alba* L. leaves by deep eutectic solvents combined with ultrasonic-assisted extraction. *Ind. Crops Prod.* 120, 147–154.